Successful full-scale deployments of advanced PGPR enhanced phytoremediation systems (PEPS) for decontamination of petroleum and salt impacted soils.

Bruce Greenberg, Xiao-Dong Huang & Perry Gerwing

WEBi Waterloo Environmental Biotechnology Inc.

EARTHMASTER ENVIRONMENTAL STRATEGIES INC.

University of Waterloo

Partners: Talisman; PennWest; Canadian Forest Oil; Devon; Baytex; ConocoPhillips; Solaction; Shell; Seaway Energy Services; Questerre Beaver River; Cenovus; Imperial Oil; URS; NorthWind; SNC Lavalin; MWH; NSERC

Outline

1. Advantages of phytoremediation
2. Overview of our phytoremediation system
3. Field tests of our phytoremediation system for petroleum and salt remediation

Examples of Remediation Methods

- Dig and dump - Any contaminant type - **$100-500/m^3**
- Soil incineration - On or off site - Organic contamination - **$200-600/m^3**
- Chemical extraction - Any type of contamination - **$250/m^3**
- Electrokinetic separation - Metals/Salts - **$200/m^3**
- Soil flushing/fracturing - Any contaminant type - **$250/m^3**
- Land farming - Natural attenuation - Small organics - **$50/m^3**
- Bioremediation - Organics - **$100/m^3**
- Phytoremediation - Any contaminant type - **$25-50/m^3**

Process of Phytoremediation



- Volatilization
- Phytodegradation
- Chelation/compartment in leaves

Salt

- Translocation: root symplast→ xylem
- Chelation/compartment in roots
- Plant uptake soil→root
- Rhizosphere Processes
- Bioavailability particle→water

Rhizodegradation - PHC

Advantages of Phytoremediation

1. Improves the quality of soil
2. It is driven by solar energy and suitable to most regions and climates
3. It is cost effective and technically feasible
4. **Plants** provide **sufficient biomass** for rapid remediation; **promote high rhizosphere activity**
5. Restoration in a reasonable time frame - 2 to 3 years
6. Can be used effectively at **remote sites**
7. Effective for remediation of PHC and salt – relevant to the energy industry

Development, Proof, and Application of PGPR Enhanced Phytoremediation Systems (PEPS)

Over 10 years of research with full-scale field studies at each stage of development and application

1. **PHC:** sites in AB, BC, QC, and ON (2004-10)
2. **Gas station:** site fully remediated in 1 summer (2007)
3. **Salt:** sites in SK, AB and NWT (2007-10)

EARTHMASTER ENVIRONMENTAL STRATEGIES INC.

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Description of the PGPR Enhanced Phytoremediation System (PEPS)

Physical soil treatment: Till the soil: exposure to sunlight and air
Exposure to sunlight photooxidizes contaminants

Bioremediation: Inoculation of PAH/TPH degrading bacteria (generally skipped in the field → already present)

Phytoremediation: Growth of plants <u>with PGPR</u>

- **<u>PGPR:</u> Plant growth promoting rhizobacteria.**
- **Prevent the synthesis of stress ethylene.**
- **<u>PGPR</u> are applied to the seeds prior to sowing → NOT Bioaugmentation**
- **Grass species used generally**
- **Effect depth of remediation ~ 0.5 m**

Interaction of a PGPR Containing ACC Deaminase with a Plant Seed or Root



Plant growth promoting rhizobacteria (PGPR)

Natural, non-pathogenic strains of PGPR (usually *Pseudomonads*)

We have isolated PGPRs from ON, AB, SK and the NWT

PGPR are applied to seeds prior to planting

Research and Development of the PEPS for PHC Remediation

1. **Sarnia, ON – land farm – 4 year study**

   Oil sludge – PHC contaminated soil

   (15% w/w – 60% F3 (C16-C34), 30% F4 (C34 – C50)

2. Turner Valley, AB – 3 year study

3. Hinton, AB – 2 year study

Waterloo Environmental Biotechnology Inc.

Sarnia, ON – Land Farm

- Planted barley/fescue/ryegrass
- Plants were treated with PGPR (UW3 and UW4) using a mechanical seed treater
- PHC remediation from:
    - 15% - 3% in 4 years with PEPS
    - 15% - 8% in 4 years for plants w/o PGPR
    - 15% - 11% in 4 years w/o plants



Conclusions on Development of the PEPS

- 100% increase in plant biomass due to PGPR, root growth to 50 cm below ground level
- 30 to 40% remediation per year with PEPS; 100% faster than plants without PGPR
- Rhizosphere microbes (esp. PHC degraders) elevated 10 to 100 fold with the PEPS - microbes and plants consume PHC
- Very low ^{14}C detected in soil microbial fatty acids – Carbon came from PHC metabolism (PHC has no ^{14}C)
- Very low ^{14}C in CO_2 that evolves from soil – PHC has been mineralized to CO_2
- No PHC detected in plant tissue as it disappears from the soil
- CCME PHC analytical method used effectively to show extent of remediation

EARTHMASTER ENVIRONMENTAL STRATEGIES INC.

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Phytoremediation of PHC

(A) Bioavailability of PHC

(B) General processes affecting rhizoremediation

(C) Microbial aerobic PHC degradation – rhizosphere supported by plants

(D) Possible microbial oxygenation pathway of PHC to form a fatty acid









Application of PEPS for PHC Remediation – 1st Generation Full Scale Sites for Proof of Concept (2007- 09)

All sites planted with oats, tall fescue and ryegrass treated with PGPR – All sites met applicable criteria

1. Hinton 2, AB – Complete remediation in 2 years – Diesel invert drilling waste
2. **Edson, AB –** Complete remediation in 2 years – Diesel invert drilling waste
3. Peace River, AB – Complete remediation in 3 years – Flare pit material
4. Steinbach, MB – Complete remediation in 1 year – Gas station site
5. **Quebec City, QC –** Tier 1 criteria met in one year

EARTHMASTER ENVIRONMENTAL STRATEGIES INC.

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Edson, AB – Site and Sampling Map (2008)

Soil Impact – PHC (Diesel Invert; 85% F3)



Area:
120 m x 100 m
= 12,000 m^2
= 3 acres

EARTHMASTER ENVIRONMENTAL STRATEGIES INC.

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Edson, AB – Beginning and Mid-Season (2008)

Soil Impact – PHC (Diesel Invert; 85% F3)



EARTHMASTER ENVIRONMENTAL STRATEGIES INC.

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Edson, AB – PHC Remediation (2007- 08)

Soil Impact – PHC (Diesel Invert; 85% F3)

★ **In June 2007, 9 of 13 sampling points above Tier 1 criteria (F3 > 1300 mg/kg)**



EARTHMASTER ENVIRONMENTAL STRATEGIES INC.

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Gravimetric Total PHC vs Analytical Lab F3



- Can follow F3 remediation with a less expensive method
- Final remediation is confirmed using accredited lab

Edson, AB – PHC Remediation (2007- 08)

Soil Impact – PHC (Diesel Invert; 85% F3)



- **Remediation goals were met**
- **No points over Alberta Tier 1 criteria**

EARTHMASTER ENVIRONMENTAL STRATEGIES INC.

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Quebec City, QC – End of Season (2009)

Soil Impact – PHC

Quebec City, QC – End of Season (2009)

Soil Impact – PHC



Quebec City, QC – End of Season (2009)

Soil Impact – PHC



Second Generation Full Scale Sites

1. Three sites near Dawson Creek, BC
2. One site near Swan Hills, AB
3. One site near Hinton, AB
4. One site near Edson, AB
5. One site near Red Earth Creek, AB
6. One site in Northern BC

All sites worked with very similar results
All sites planted with tall fescue, ryegrass, and/or oats treated with PGPR.

Northern BC near NWT Border – Mid/End of Season (2010)

Soil Impact – PHC (Diesel Invert)



Mid-Season

End of Season

Northern BC near NWT Border – Site Map (2010)

Soil Impact – PHC (Diesel Invert)



Site Area:

- 2400 m^2
- 0.6 acres

Northern BC – EPH Remediation (2010)



$EPH_{C10-C19}$ Remediation (Maxxam)

- 6 out of 8 sampling points showed a decrease in $EPH_{C10-C19}$ levels over two months
- All sampling points had $EPH_{C10-C19}$ level above 1000 mg/kg criteria
- At the end of the season the average **$EPH_{C10-C19}$ level decreased by 29%**, from 3659 mg/kg in July to 2608 mg/kg in September



$EPH_{C19-C32}$ Remediation (Maxxam)

- 6 out of 8 sampling points showed a decrease in $EPH_{C19-C32}$ levels over two months
- Only 3 out of 9 sampling points had $EPH_{C19-C32}$ level above 1000 mg/kg criteria in September
- At the end of the season the average **$EPH_{C19-C32}$ level decreased by 27%**, from 1335 mg/kg in July to 979 mg/kg in September

Swan Hills, AB – End of Season (2009 – 10)

Soil Impact – PHC





Planted un-impacted plot

Soil sampling point

EARTHMASTER ENVIRONMENTAL STRATEGIES INC.

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Swan Hills, AB – End of Season (2009 – 10)

Soil Impact – PHC

Impacted Plot



Un-impacted Plot

EARTHMASTER ENVIRONMENTAL STRATEGIES INC.

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Swan Hills, AB – End of Season (2009 – 10)

Soil Impact – PHC

Impacted Plo

Sample date	Soil-type	Plant Biomass (g/m^2) ± SD
October 2009	Impacted	427.7 ± 88.06
October 2009	Un-impacted	438.1 ± 53.56



Un-impacted Plot

EARTHMASTER ENVIRONMENTAL STRATEGIES INC.

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Swan Hills, AB – F2/F3 Results





F2 & F3 analysis performed by Maxxam. Method used: cold shake extraction, single silica-column clean-up

EARTHMASTER ENVIRONMENTAL STRATEGIES INC.

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Phytoremediation Cost analysis for a typical PHC impacted site - Edson

- Collaborative project between Earthmaster Environmental and Waterloo Environmental Biotechnology
- Volume of impacted material – 460 m^3 of diesel invert drilling mud was originally spread over 1.07 ha
- 1.07 ha impacted to a depth of 0.3 m or <u>3,210 m^3 of PHC impacted material</u>
- The costs for the entire project was: $104,000 or **$32.50/m^3**
  - Includes all Earthmaster, WEBi and 3rd party costs
- Landfilling this material would have cost **$80/$m^3$**
  - Assumes a 2 h truck turnaround time
  - No backfill required – if backfill was required the cost would rise to **$90/$m^3$**

EARTHMASTER ENVIRONMENTAL

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Conclusions for PHC Remediation

SUCCESS

- Achieved PHC remediation: 4 sites brought to closure, 6 second generation sites progressing well towards closure

PERFORMANCE PREDICTIONS FOR PEPS

- PHC up to 10,000 mg/kg – Remediation in 2 to 4 years
- PHC > 10,000 mg/kg – Remediation in 3 to 6 years
- In all cases, soil becomes non-toxic before regulatory criteria are met – Site specific risk assessment possible

COST

- Actual cost for the Edson site (3,400 m^3) was \$33/m^3
- Cost to landfill (landfill 1 h from site) would have been > \$80/m^3
- Cost differential increases as site become more remote or volume of impacted material increases

EARTHMASTER ENVIRONMENTAL

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Development of PEPS for Salt Impacted Sites

EARTHMASTER ENVIRONMENTAL STRATEGIES INC.

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Plant responses to salinity

- Inhibited germination
- Decreased water uptake → Low water potential (drought)
- Unbalanced sodium/potassium ratios
- Inhibition of photosynthesis
- Increased reactive oxygen species (ROS)
- Increased **ethylene production**

Salinity effects mostly negligible (or salt deprived)	Yields of very sensitive crops may be restricted	Yields of many crops diminished	Only tolerant plants grow	Only a few very tolerant plants can grow

0 2 4 8 16

EC_e (dS/m)

EARTHMASTER ENVIRONMENTAL STRATEGIES INC.

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Sites for Development of PEPS for Salt Remediation

1. Cannington Manor, SK
2. Alameda, SK
3. Kindersley, SK
4. Brazeau, AB
5. Norman Wells, NWT
6. Weyburn, SK (7 sites)
7. Provost, AB
8. Red Earth, AB

EARTHMASTER ENVIRONMENTAL STRATEGIES INC.

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Lab Research Summary of PEPS for Salt Impacted Soils

- 50 to 100% increases in plant growth due to PGPR with root growth to 50 cm
- Plants can grow on soils with EC_e ~ 25 dS/m
- ON, SK, and NWT PGPRs all worked well
- PGPRs protected against inhibition of photosynthesis and plant membrane damage
- Levels of salt up-take to plant foliage: 50 – 75 g NaCl/kg dry weight
- Remediation can be based on up take of salt into foliage
- Phytoremediation is feasible for soils with EC_e of 15 to 20 dS/m in about 5 years

EARTHMASTER ENVIRONMENTAL

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Norman Wells, NWT – End of Season (2010)

Soil Impact – Salt

Norman Wells, NWT – End of Season (2008)

Soil Impact – Salt



Plants used: slender wheatgrass and red fescue

No soil conditioning

Norman Wells, NWT – End of Season (2009)

Soil Impact – Salt



Plants used: slender wheatgrass, ryegrass and red fescue

Soil conditioned

Norman Wells, NWT – End of Season (2010)

Soil Impact – Salt



Norman Wells, NWT – End of Season (2010)

S



Salt Remediation





Norman Wells, NWT – End of Season (2009)

Soil Impact – Salt

High salt plant material was mowed and removed from the site

Norman Wells, NWT – End of Season (2009)

Soil Impact – Salt



Plant Biomass (dry wt g/m^2 SE)						
Year	**Plot A**		**Plot B**		**Plot C**	
2009	300	26	397	50	623	44
2010	393	16	592	40	525	20

High salt plant material was mowed and removed from the site

Weyburn, SK – 1 of 7 sites: Soil Salinity (EC_e) Map (2010, Year 1)

Soil Impact – Salt



Surface soil EC_e at each sample point

Field Averages:
= **8.35 dS/m (0-60 cm)**

= 5.67 dS/m (0-15 cm)
= 7.48 dS/m (15-30 cm)
= 11.89 dS/m (30-60 cm)

Site Area:
- 4500 m^2
- 1.1 acres

EARTHMASTER ENVIRONMENTAL STRATEGIES INC.

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Weyburn, SK – 1: End of Season (2010, Year 1)

Soil Impact – Salt



EARTHMASTER ENVIRONMENTAL STRATEGIES INC.

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Weyburn, SK – 1: End of Season (2010, Year 1)

Soil Impact – Salt



Weyburn, SK – 2 of 7 sites: Soil Salinity (EC_e) Map (2010, Year 1)

Soil Impact – Salt

Site Area:

- 12 acres

Field Averages:

= **4.61 dS/m (0-60)**

= 3.25 dS/m (0-15 cm)

= 3.43 dS/m (15-30 cm)

= 3.82 dS/m (30-60 cm)



EARTHMASTER ENVIRONMENTAL STRATEGIES INC.

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Weyburn – 2, SK: End of Season (2010, Year 1)

Soil Impact – Salt



EARTHMASTER ENVIRONMENTAL STRATEGIES INC.

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Weyburn – 2, SK: End of Season (2010, Year 1)

Soil Impact – Salt





EARTHMASTER ENVIRONMENTAL STRATEGIES INC.

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Provost, AB: Mid-Season (2009, Year 1)

Soil Impact – Salt



High salt hot spots and poor soil from pipeline construction:
EC_e: 13-17 dS/m

Waterloo Environmental Biotechnology Inc.

Provost, AB: End of Season (2009, Year 1)

Soil Impact – Salt

- High salt spots have filled in with plants

- EC_e (2009): decreased from 13-17 to 4-12 dS/m
- EC_e (Spring 2010): all sampling points were below applicable targets
- **Successful remediation was achieved in 1 year**

Waterloo Environmental Biotechnology Inc.

Red Earth, AB: 2009 - 2010

Soil Impact – Salt and Herbicide (Arsenal®)

Great improvement in plant growth, therefore, can remediate herbicides

Why Use Phytoremediation?

- It works for PHC and salt remediation.
- Remediation at all sites (> 20) successful.
- Costs of PHC and salt remediation will be similar.
- Unit cost drops as the volume of material increases.
- Phytoremediation costs (all in) < half the cost of landfilling.
- Liability is reduced, not transferred to a landfill.
- Costs are spread over more than one year (2 to 4 yrs).
- The price differential relative to landfilling increases when sites become more remote.
- Purchase of backfill not required. Soils are reused.
- Tier 2 approach - if required only marginal cost increase.
- Green technology: Good PR and environmentally friendly.

EARTHMASTER ENVIRONMENTAL

WATERLOO ENVIRONMENTAL BIOTECHNOLOGY INC.

Colleagues and Partners

- **The people that do all the work**
  - Karen Gerhardt
  - Jola Gurska
  - Xiao-Ming Yu
  - Wenxi Wang
  - Mark Lampi
  - Shan Shan Wu
  - Julie Nykamp
  - Nicole Knezevich
  - Greg MacNeill
  - Xiaobo Lu
  - Scott Liddycoat
  - Han Zheng
  - Brianne McCallum
  - Jing Ma
  - Peter Mosley
  - Conrad Neufeld
  - Xiao-Dong Huang

- **Collaborators**
  - Perry Gerwing, Earthmaster

- **Partners**
  - S Willets, O Mrklas, C Gordey, ConocoPhillips Canada
  - B Moore, Devon
  - G Millard, Shell
  - J Budziak, Seaway
  - P Coldham, Questerre Beaver River
  - E Harrison, Cenovus
  - L Lawlor, Imperial Oil
  - K Cryer, M Metzger, S Brown, C Chattaway, Earthmaster
  - D McMillan, SNC
  - G Stephenson, Stantec
  - T Chidlaw, MWH
  - S Steed, NorthWind
  - A Traverse, Baytex
  - G Adams, URS
  - B Chubb, Maxxam

WEBI
Waterloo Environmental Biotechnology Inc.