

**Disposition of Radioactivity 24 Hours Following Last of 10 Daily Oral Gavage
Administrations of 20 mg/kg [¹⁴C]Tetrabromobisphenol A
to Male F-344 Rats (Group 6)**

Disposition Summary^a
% Dose Recovered in Excreta and Tissues (24 hours)

| Rat Identification number | 50315-01 | 50315-02 | 50315-03 | 50315-04 | Mean | SD ^b |
|------------------------------|-----------------|----------|----------|----------|--------------|-----------------|
| Excreta | - | - | - | - | - | - |
| Feces | 97.96 | 96.76 | 97.54 | 99.43 | 97.9 | 1.12 |
| Urine | 0.3 | 0.31 | 0.27 | 0.52 | 0.35 | 0.11 |
| Cage rinse | 0.07 | 0.12 | 0.15 | 0.12 | 0.11 | 0.03 |
| Tissues | - | - | - | - | - | - |
| Blood | ND ^c | ND | ND | ND | ND | - |
| Cecum and intestinal content | 0.85 | 1.42 | 0.66 | 0.55 | 0.87 | 0.38 |
| Extra-GI Tissues | 0.07 | 0.09 | 0.17 | 0.08 | 0.10 | 0.04 |
| Total recovery | 99.25 | 98.7 | 98.79 | 100.7 | 99.36 | 0.92 |

^a The target dose was 20 mg/kg body weight (50 µCi/kg). Rats received a daily dose of 20 mg/kg body weight for 10 days. Excreta values are cumulative.

^b SD = standard deviation (SD). Shown is the mean ± standard deviation for the group (N = 4).

^c ND = not detectable.