

**Experiment Number:** C16008-01

**Test Type:** TOX

**Route:** Dosing in Bedding

**Species/Strain:** Mouse/B6C3F1/N

**I06: Mean Feed Consumption**

**Test Compound:** Crumbrubber various

**CAS Number:** CRUMBRUBBERVARIOUS

**Date Report Requested:** 02/12/2018

**Time Report Requested:** 10:43:27

**Lab:** Battelle

**C Number:**

C16008-01

**Cage Range:**

All

**Date Range:**

All

**Reasons For Removal:**

All

**Removal Date Range:**

All

**Treatment Groups:**

All

**Study Gender:**

Female

**Experiment Number:** C16008-01  
**Test Type:** TOX  
**Route:** Dosing in Bedding  
**Species/Strain:** Mouse/B6C3F1/N

**I06: Mean Feed Consumption**  
**Test Compound:** Crumbrubber various  
**CAS Number:** CRUMBRUBBERVARIOUS

**Date Report Requested:** 02/12/2018  
**Time Report Requested:** 10:43:27  
**Lab:** Battelle

Females								
Phase	Litter ID	Days	Treatment Groups (wt./wt.)					
			0:0			50:50		
			Wt (g/animal/day)	Wt (g/kg/animal/day)	N	Wt (g/animal/day)	Wt (g/kg/animal/day)	N
Study		0 - 4	3.6 ± 0.0	174.6 ± 2.5	15	3.8 ± 0.1	183.8 ± 4.5	15
		4 - 7	2.6 ± 0.2	125.4 ± 9.2	15	3.2 ± 0.2 **	153.8 ± 7.8 **	15
		7 - 11	3.5 ± 0.1	161.8 ± 3.3	15	3.7 ± 0.1	173.1 ± 5.9	15
		11 - 14	2.9 ± 0.0	137.2 ± 2.7	10	3.3 ± 0.0 **	155.9 ± 1.6 **	10
		0 - 14	3.2 ± 0.0	154.6 ± 3.3	10	3.6 ± 0.1	170.3 ± 3.4 **	10

**Experiment Number:** C16008-01

**Test Type:** TOX

**Route:** Dosing in Bedding

**Species/Strain:** Mouse/B6C3F1/N

**I06: Mean Feed Consumption**

**Test Compound:** Crumbrubber various

**CAS Number:** CRUMBRUBBERVARIOUS

**Date Report Requested:** 02/12/2018

**Time Report Requested:** 10:43:27

**Lab:** Battelle

#### LEGEND

---

Data are displayed as mean  $\pm$  SEM

N is the number of animals (excluding unweaned pups)

Statistical analysis performed by Jonckheere (trend) and Shirley or Dunn (pairwise) tests (unless otherwise noted).

Statistical significance for the control group indicates a significant trend test

Statistical significance for a treatment group indicates a significant pairwise test compared to the vehicle control group

\* Statistically significant at  $P \leq 0.05$

\*\* Statistically significant at  $P \leq 0.01$

Consumption is not reported for the non-pregnant animals during gestation and lactation phases

Consumption is not reported for animals during mating

Decrease in N for the F0 Females data for LD 10 to 13 was because 1 value in the 3000 ppm group was an outlier.

Statistical analysis performed using the Mann-Whitney U test for comparing the two groups.

**\*\* END OF REPORT \*\***