

Bioavailability Changes in Sediments and Bioaccumulation in Fish

Changes in uptake of polychlorinated biphenyls (PCBs) in fish after remediation of their aquatic environment may be predicted, according to researchers at the University of Maryland, Baltimore County. They measured freely dissolved concentrations of PCBs in water and applied mathematical models to predict the effectiveness of sediment remediation. The study is one step toward understanding how PCB bioavailability changes in sediment as a result of activated carbon amendments, a method to sequester PCBs, can influence transfer of PCBs to fish.

PCBs are a family of chemical compounds that were widely used in industrial applications and still persist in the environment in air, water, soil and sediments, and foods. Activated carbon amendments have been demonstrated to reduce porewater concentrations of PCBs, but little is known about how these changes affect bioaccumulation in fish. Within aquatic ecosystems, PCBs accumulate up the food chain. Fish consumption is a primary source of PCB exposure in people.

Evaluating activated carbon treatment

The study, led by Upal Ghosh, Ph.D., evaluates the use of activated carbon, the same technology employed in many water filters, to bind contaminants in sediment, reducing availability in the aquatic ecosystem. Researchers performed laboratory aquarium experiments and compared their results to model predictions to explore how PCBs affect exposure pathways and bioaccumulation in fish.

They measured PCBs using passive sampling, an effective method for measuring freely dissolved concentrations in water, and found that PCBs were reduced by more than 95 percent with activated carbon. They also saw a 99 percent reduction in PCB concentrations in sediment porewater. They measured PCBs in fish tissue lipids and found that the amendment reduced PCB uptake in fish by 87 percent after 90 days. These results show that bioavailability changes in sediment are reflected in uptake in fish.



SRP trainees Hilda Fadaei (UMBC) and Aaron Watson (Institute of Marine and Environmental Technology) measure PCB levels in water and fish tissue in aquarium experiments. (Photo courtesy of Upal Ghosh)

The researchers then tested two existing aquatic bioaccumulation models that may predict PCB uptake changes after sediment is treated with activated carbon. While the equilibrium and kinetic models are available, there is limited knowledge on their ability to predict levels in fish after treatment of sediment to reduce PCB exposure.

They found that the kinetic model was a better predictor and generally within a factor of two for total PCBs measured in fish. Based on these results, the researchers concluded that this bioaccumulation model can be linked to other existing fate and transport models to predict effectiveness of sediment remediation in reducing PCB uptake in fish over time. After validation, these types of models can also be used as a reliable means of back-calculating the activated carbon amendment dose required to optimize the desired end point in recovery of fish tissue PCB levels.

Translating the research into the field

This study also helps show the effectiveness of remediation using activated carbon and may spur translation of the technology to the field. SediMite, an activated carbon amendment technology developed by Ghosh, is currently being used in the remediation of sediments contaminated with PCBs. The product packages activated carbon with clay and sand into pellets that are convenient to handle and deliver. It slowly breaks up over time and mixes through natural processes when added to the water column.

A project at Mirror Lake in Dover, Delaware is the first full-scale implementation of activated carbon amendment into sediments. Mirror Lake and downstream areas have been in decline for several decades from stormwater runoff contamination from chemicals, excess nutrients, bacteria, and invasive plant species. If left untouched and with no further contamination, the Delaware Department of Natural Resources and Environmental Control forecasts it would take several decades for the lake to clean itself naturally and for fish in the lake to be safe to eat. With the technology, DNREC scientists anticipate a reduction of PCB contaminants in fish tissue to below consumption advisory levels within a few years.

The U.S. Environmental Protection Agency, in its recent notice of action for the Housatonic River Superfund Site, has also asked that activated carbon amendment to sediment be considered as part of the final remedy. This study helps to show how activated carbon affects PCB uptake in fish and how researchers can make reasonable assessments of long-term recovery of fish stocks.

For more information, contact:

Upal Ghosh

University of Maryland, Baltimore County
Department of Chemical, Biochemical, and Environmental Engineering
1000 Hilltop Circle
Baltimore, MD 21250
Tel: 410-455-8665
Email: ughosh@umbc.edu

To learn more about this research, please refer to the following source:

Fadaei H, Watson A, Place A, Connolly J, Ghosh U. 2015. Effect of PCB bioavailability changes in sediments on bioaccumulation in fish. *Environ Sci & Technol* 49(20):12405-12413. doi: [10.1021/acs.est.5b03107](https://doi.org/10.1021/acs.est.5b03107)

